

A THEOREM ON APPROXIMATION BY ALGEBRAIC POLYNOMIALS OF FUNCTIONS WITH GIVEN GENERALISED MODULUS OF SMOOTHNESS

M. K. POTAPOV AND F. M. BERISHA

There are well known theorems on approximation by algebraic polynomials of functions, where the generalised modulus of smoothness is defined by means of symmetric operator of generalised translation. It is of interest obtaining the same results for a modulus of smoothness defined by an asymmetric operator of generalised translation.

We introduce such an operator, define the generalised modulus of smoothness by its means, and obtain the direct and inverse theorems in approximation theory for it.

Denote by $L_{p,\alpha}$, $1 \leq p \leq \infty$, the set of functions f such that $f(x)(1-x^2)^\alpha \in L_p$, and put

$$\|f\|_{p,\alpha} = \left\| f(x)(1-x^2)^\alpha \right\|_p.$$

By $E_n(f)_{p,\alpha}$ we denote the best approximation of a function $f \in L_{p,\alpha}$ by algebraic polynomials of degree not greater than $n-1$, in $L_{p,\alpha}$ metrics.

For a function $f \in L_{p,\alpha}$ we define the asymmetric operator of generalised translation:

$$\begin{aligned} \hat{\tau}_t(f, x) &= \frac{1}{\pi(1-x^2)\cos^4 t/2} \\ &\times \int_0^\pi \left\{ 2 \left(\sqrt{1-x^2} \cos t + x \sin t \cos \varphi + \sqrt{1-x^2}(1-\cos t) \sin^2 \varphi \right)^2 \right. \\ &\left. - 1 + \left(x \cos t - \sqrt{1-x^2} \sin t \cos \varphi \right)^2 \right\} f \left(x \cos t - \sqrt{1-x^2} \sin t \cos \varphi \right) d\varphi. \end{aligned}$$

By means of that operator of generalised translation we define the generalised modulus of smoothness by

$$\hat{\omega}(f, \delta)_{p,\alpha} = \sup_{|t| \leq \delta} \|\hat{\tau}_t(f, x) - f(x)\|_{p,\alpha}.$$

Theorem 1. *Let given numbers p and α be such that $1 \leq p \leq \infty$;*

$$\begin{aligned} 1/2 < \alpha \leq 1 & \quad \text{for } p = 1, \\ 1 - \frac{1}{2p} < \alpha < \frac{3}{2} - \frac{1}{2p} & \quad \text{for } 1 < p < \infty, \\ 1 \leq \alpha < 3/2 & \quad \text{for } p = \infty. \end{aligned}$$

Let $f \in L_{p,\alpha}$. For every natural number n the following inequalities hold

$$C_1 E_n(f)_{p,\alpha} \leq \hat{\omega}(f, 1/n)_{p,\alpha} \leq C_2 \frac{1}{n^2} \sum_{\nu=1}^n \nu E_\nu(f)_{p,\alpha},$$

where the positive constants C_1 and C_2 do not depend on f and n .

1991 *Mathematics Subject Classification.* Primary 41A35, Secondary 41A50, 42A16.

This work was done under the support of the Russian Foundation for Fundamental Scientific Research, Grant #97-01-00010 and Grant #96/97-15-96073.

М. К. РОТАРОВ, МЕХАНИКО-МАТЕМАТИЧЕСКИ ФАКУЛЬТЕТ МГУ, МОСКВА 117234, РОССИЯ

F. M. BERISHA, FACULTY OF MATHEMATICS AND SCIENCES, UNIVERSITY OF PRISHTINA,
NËNA TEREZË 5, 10000 PRISHTINË, KOSOVO
E-mail address: `faton.berisha@uni-pr.edu`